

source: <https://doi.org/10.7892/boris.123085> | downloaded: 5.5.2023

- 2) The predicted outcome of established ICU scores is mortality and not occurrence of organ dysfunction. After the initial decision to admit the patient to the ICU for full treatment, the prediction of mortality by scoring is not of great importance for further treatment decisions.
- 3) For simplicity and feasibility of routine use, only the most pathologic value in a dataset obtained during first day of ICU admission is used for scoring, whereas changes of organ function variables over time—which may be of greater prognostic relevance—are not integrated.
- 4) Organ function parameters can be influenced by the established organ support or replacement modalities and changes in treatment intensity, which are not adequately included as scoring variables.

Despite the obvious potential and the promise of supporting a wide range of medical functions, including automated patient surveillance and clinical decision support, prognostic modeling based on “big data” using modern machine learning methods is not well established in an ICU context (9). To date, only basic predictive models for the detection of deterioration of organ functions have been developed for specific disease conditions, such as hemodynamic instability (10, 11), sepsis (12), and lung failure (13). Given the complexity of the task, systems which include and integrate a multitude of patient data accumulating over time during the treatment process to predict the course of illness severity in the immediate future do not exist yet. However, the advent of large-scale healthcare information systems offers the opportunity to advance medicine to the digital age. Using this information smartly requires further extensive research and the collaboration of clinicians and data specialists as was exemplary done by Deliberato et al (4). To bring the potential of big data to the bedside, we will need to make patient data accessible for research, define clinically meaningful endpoints, and develop the right analytic methods.

## REFERENCES

1. Capuzzo M, Valpondi V, Sgarbi A, et al: Validation of severity scoring systems SAPS II and APACHE II in a single-center population. *Intensive Care Med* 2000; 26:1779–1785
2. Higgins TL, Teres D, Copes WS, et al: Assessing contemporary intensive care unit outcome: An updated Mortality Probability Admission Model (MPM0-III). *Crit Care Med* 2007; 35:827–835
3. Moreno RP, Metnitz PG, Almeida E, et al; SAPS 3 Investigators: SAPS 3—From evaluation of the patient to evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med* 2005; 31:1345–1355
4. Deliberato RO, Ko S, Komorowski M, et al: Severity of Illness Scores May Misclassify Critically Ill Obese Patients. *Crit Care Med* 2018; 46:394–400
5. Ng M, Fleming T, Robinson M, et al: Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014; 384:766–781
6. Saeed M, Villarroel M, Reisner AT, et al: Multiparameter intelligent monitoring in intensive care II: A public-access intensive care unit database. *Crit Care Med* 2011; 39:952–960
7. Saeed M, Lieu C, Raber G, et al: MIMIC II: A massive temporal ICU patient database to support research in intelligent patient monitoring. *Comput Cardiol* 2002; 29:641–644
8. Moss TJ, Lake DE, Calland JF, et al: Signatures of Subacute Potentially Catastrophic Illness in the ICU: Model Development and Validation. *Crit Care Med* 2016; 44:1639–1648
9. Pirracchio R, Petersen ML, Carone M, et al: Mortality prediction in intensive care units with the Super ICU Learner Algorithm (SICULA): A population-based study. *Lancet Respir Med* 2015; 3:42–52
10. Cao H, Eshelman L, Chbat N, et al: Predicting ICU hemodynamic instability using continuous multiparameter trends. *Conf Proc IEEE Eng Med Biol Soc* 2008; 2008:3803–3806
11. Eshelman LJ, Lee KP, Frassica JJ, et al: Development and evaluation of predictive alerts for hemodynamic instability in ICU patients. *AMIA Annu Symp Proc* 2008; 2008:379–383
12. Lindner HA, Balaban Ü, Sturm T, et al: An algorithm for systemic inflammatory response syndrome criteria-based prediction of sepsis in a polytrauma cohort. *Crit Care Med* 2016; 44:2199–2207
13. Huddar V, Rajan V, Bhattacharya S, et al: Predicting postoperative acute respiratory failure in critical care using nursing notes and physiological signals. *Conf Proc IEEE Eng Med Biol Soc* 2014; 2014:2702–2705